



OOA HOLDING ORDERS FOR ANTICOAGULANT MEDICATIONS FOR ALL PROCEDURES AND SURGERIES EXCEPT SPINES

Patient Name: _____

Diagnosis for Anticoagulant (Include dosage and frequency): _____

MRN: _____

Other (NSAID, HERBALS, ASA):

Surgery Date: _____

Anticoagulant: _____

Please check the appropriate hold order for the anticoagulant medication(s) listed.

AGGRENOX (ASPIRIN/DYPYRIDAMOLE) – Hold for 24-hours prior to surgery.

BRILINTA (TICAGRELOR) – Hold for 7 days prior to surgery.

ELIQUIS (APIXABAN) – Hold for 72 hours prior to surgery.

EFFIENT (PRASUGREL) – Hold for 7 days before surgery.

PLAVIX (CLOPIDOGREL) – Hold for 7 days prior to surgery.

PRADAXA (DABIGATRAN) – Hold for 5 days prior to surgery.

WARFARIN (COUMADIN) - Holding instructions will be defaulted to patient’s anticoagulation Clinic (ACC) or prescribing provider, notification must be sent to respective ACC with patient demographics, surgery type, and date for surgery.

XARELTO (RIVAROAXABAN) – Hold for 72 hours prior to surgery.

ASPIRIN - Please continue ASA uninterrupted at the time of surgery if prescribed by your health care provider, otherwise discontinue 14 days prior to surgery.

VITAMINS/SUPPLEMENTS/HERBALS- Stop ALL herbal supplements 10 days prior to surgery.

If hold order differs from any of the above, please indicate below:

Include medication(s) and how long to hold the medication(s).

Please check the appropriate restart of anticoagulant therapy post-surgery.

12-hours post-surgery

24-hours post-surgery

48-hours post-surgery

Other: _____

Provider Signature: _____ Date: _____