

*Life in Motion*

**Dr. Hammer**  
3909 9<sup>TH</sup> AVE SW  
Olympia, WA 98502  
360-570-3460

**Discharge Instructions**  
**Posterior Cervical Fusion**

**Medications:**

Avoid taking Advil, Motrin, Ibuprofen, Aleve, or other NSAIDs (non-steroidal anti-inflammatory drugs) for 3 months after surgery.

Avoid taking blood thinners such as Aspirin, Plavix, Coumadin, Warfarin, or Argatroban **UNLESS EXPLICITLY CLEARED BY DR. HAMMER.**

You will be given a list of medications for home. This will include pain medication, a stool softener and Miralax.


You can take Tylenol for pain if you do not need narcotic pain medication, but remember that many kinds of narcotics, such as Vicodin, have Tylenol in them. Avoid taking more than 3000 mg of Tylenol (acetaminophen) in a 24 hour period

You should resume your previous medications. Any changes will be noted on your discharge medication list.

**\*\*CONSTIPATION:** The pain medication often causes constipation, take the prescribed stool softener (Colace) and Miralax daily as long as you are on the pain medication. You may also add Senna to Miralax. Limiting the amount of narcotics will help improve constipation. Increase fluids, fresh fruit (prunes) and vegetables high in fiber. If you have not had a bowel movement within 1-2 days after returning home from the hospital, over the counter medication may be purchased at most drug stores: Magnesium Citrate (liquid), suppository, or enema. Start this by day 3 or 4 if unable to have a bowel movement. Please call our office for further suggestions to help with this.

**Neck Incision:**

Your neck wound will have a dressing that is made up of aquacel dressing. The dressing is not waterproof and must not get wet. You may remove it on post op day #4 to shower. There are staples under the dressing that will be removed at your post op appointment. These help hold the skin together until it heals. You will be contacted and scheduled by Dr. Halpin's office for a wound check appointment for 10-14 days after surgery. Any staples or residual steri-strips will be removed at that time.



## *Life in Motion*

**Call the office at (360) 709-6223 if there is excess drainage on the steri-strip. It is normal to see some drainage on the steri-strips, but they should not be soaked with blood.**

### **Collar:**

You may have been asked by Dr. Hammer to wear a cervical collar. The purpose is to remind you not to move your neck, but it will not keep you from moving your neck if you try. Unless specified by Dr. Hammer, you do not need to wear your cervical collar in bed when your head is on a pillow, while sitting up in a chair or while eating. You can take it off to shower. You should wear the cervical collar through your follow up visit. You may discuss the duration you need to wear the collar at that time but most people will be taken out of the collar at 6 weeks.

The collar does not come with replacement pads. You may wash them in the washer. A good time to do this is while you are sitting in a chair. And not needing to wear the collar.

### **Driving:**

It is best not to drive for the first 3 weeks after surgery. This is because driving involves a lot of bending, twisting, and turning of the neck. You should not drive while you are taking narcotic pain medication.

### **Watch for the following signs and symptoms and call the office if they occur:**

Temperature greater than 101° F

Persistent drainage from the wound of blood or clear fluid

Any drainage of pus from the wound

Increasing redness and swelling around the incision

Separation of the wound edges

Do not apply any lotions, ointments, or other products to your incision unless directed by your physician

### **Fever:**

Some patients have low grade fevers (99°-101°F) in the immediate post operative period. This is almost always due to the fact that your lungs are not fully re-expanded after the general anesthetic.

You can treat this fever with Tylenol but it is important to try to cough the phlegm out of your lungs, take deep breaths to re-expand your lungs, and begin to increase your activities as tolerated

Wound infections are rare in the first couple days after surgery. If you have a low grade fever, try coughing and deep breathing.

### **Activities and Restrictions:**

Avoid heavy lifting (greater than 10 lbs), bending, or straining.

Do not sit for long periods of time, including long car rides

Avoid any activities that put a strain on your neck



## *Life in Motion*

It is normal to feel tired after the operation. Your body is using all of its energy to heal itself and therefore, it is normal to be fatigued after even small activities. Your strength will gradually improve on a daily basis. The wound will not open with activity or normal movement. Activity may cause some discomfort but it is not dangerous. Do not worry about causing harm to yourself by increasing your activities. The guiding principle for your activities should be that "if it hurts, don't do it." You can gradually increase your activities by walking short distances at first then increasing the distance before doing anything else.

You should avoid returning to work until cleared by Dr. Halpin

### **Showering:**

You may shower on the 4<sup>th</sup> day after surgery. Remove the bandage as it is not waterproof and a wet bandage may cause an infection. **Do not scrub the incision.** Let the water run over it and pat it dry. Do not submerge the incision underwater in a bathtub, swimming pool, or hot tub until the wound has completely healed.

### **Diet:**

You may resume your regular diet. Nutrition is extremely important for wound healing. You may be instructed to take vitamin supplements such as a multivitamin, and Calcium with Vitamin D. These can be purchased at most grocery stores or pharmacies. In addition, nutritional shakes such as Ensure or Boost can also provide supplemental nutrition.

### **Symptoms:**

You may experience pain, numbness, weakness, or tingling that was present prior to surgery that persists. This may last awhile as the nerve needs time to heal.

### **General:**

Smoking inhibits wound healing and increases your chances of the bone fusion not healing, creating a "pseudoarthrosis." This can lead to persistent pain and require additional surgery in the future. **Do not smoke!**

Remember; ask your nurse or physician if you have any questions or if you do not understand any instructions. You may also call the clinic with any questions.

**For any post-operative problems or concerns, call  
(360) 709-6223**